



Acupuncture for cosmetic use: a systematic review of prospective studies

Byung–Cheul Shin, KMD, PhD^{1,2}, Kyeong–Tae Lim, KMD²

¹Division of Clinical Medicine, School of Korean Medicine, Pusan National University, Yangsan, Rep. of Korea, ²Department of Korean Rehabilitation Medicine, Pusan National University Korean Medicine Hospital, Yangsan, Rep. of Korea

Background: Despite the use of cosmetic acupuncture in clinical fields, there is little evidence regarding its effects and safety for cosmetic use.

Objective: Objectives of this article are to review the literature and systematically evaluate the effects and current use of acupuncture for cosmetic purposes.

Methods: We searched 15 English, Chinese, Japanese, and Korean databases by using relevant keywords. All prospective studies on acupuncture for cosmetic use were considered; we conducted literature analyses to determine the current status of cosmetic acupuncture use and extracted data related to its effects.

Results: Two randomized controlled trials and 5 single–armed prospective trials with 216 participants (150 from randomized controlled trials, 66 from single–armed studies) were included. All 7 studies reported positive results for at least one of the following: total effective rate (n=2), squares and counts of wrinkles (n=1), facial blood circulation (n=1), facial size reduction and improvement of skin condition (n=1), anti–aging by a patient–oriented self–assessment of facial elasticity scale (n=1), and water and oil content of facial skin (n=1). Four studies reported itching, dry desquamation, bruising, bleeding, pain, and swelling as adverse events. Six of 7 trials reported statistical flaws, and specific information regarding acupuncture rationale (42.9%), other components of treatment (57.1%), and practitioner background (0%) were rarely reported. However, details regarding patient indications for treatment, treatment regimen, and control interventions were consistently reported (100%) in the Standards for Reporting Interventions in Clinical Trials of Acupuncture (STRICTA) reporting guideline.

Conclusion: This systematic review suggests that a few trials regarding acupuncture for cosmetic use have incorporated rigorous designs. Although the studies reported generally positive results with tolerable safety, the methodology of the trials should be updated in a rigorous manner, including the use of randomized, sham–controlled studies with standardized interventions, in order to provide sufficient evidence regarding the effects of cosmetic acupuncture.

Keywords: acupuncture; cosmetic use; miso facial acupuncture; prospective study; systematic review

Introduction

In recent years, the popularity of cosmetic procedures has increased, concomitant with extended life expectancy. According to global statistics released by the International Society of

Aesthetic Plastic Surgery in 2017, there were 13,209,539 non-surgical cosmetic procedures in 2016, approximately 10% more than the number performed in 2015. In both 2015 and 2016, the number of non-surgical procedures exceeded those of total surgical procedures. Interest in cosmetic acupuncture is similarly

Received October 11, 2018; Accepted November 26, 2018

Corresponding author: Byung–Cheul Shin

E–mail: drshinbc@pusan.ac.kr, ORCID: <http://orcid.org/0000-0002-0059-2689>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non–Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted non–commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2018 Korean Society of Korean Cosmetic Surgery and Medicine (KSKCS & KCCS).

increasing [1,2].

Cosmetic acupuncture is a type of cosmetic procedure, defined as the use of acupuncture on the head, face, and neck for cosmetic purposes without surgery and injection, such as to enhance facial muscle and skin tone, or for moisturizing purposes [3,4]. The primary aims of cosmetic acupuncture in Korea include facial wrinkle care, anti-aging, and reduction of facial size [5]. Though acupuncture has a long history, spreading from ancient China to Korea, Japan, and Western society in the early 1970s, acupuncture for cosmetic use is a recent concept resulting from the sharp increase in economic status at the end of the 20th century [4].

Despite the use of cosmetic acupuncture in clinical fields, there are few reports on the evidence of its effects and safety for cosmetic use [4,5]. There have been several reports regarding the use of cosmetic acupuncture for the rejuvenation of facial skin [5,6] but these did not constitute systematic analyses of its effects or safety. Recently, an article was published analyzing facial cosmetic acupuncture in Korea [5], but it did not constitute a systematic review.

Previously published reviews did not adhere to Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) reporting guidelines [7], and were not conducted systematically. Therefore, the aims of this study were to review the literature to determine the current status of the use of facial cosmetic acupuncture and to systematically evaluate the effects and safety of cosmetic acupuncture while adhering to the PRISMA reporting guidelines.

Materials and methods

Data sources and searches

The following electronic databases were searched up to August 2018. We searched 4 English databases (PubMed, Ovid LWW Medline, EMBASE, and Cochrane Library), 1 Chinese database (China National Knowledge Infrastructure [CNKI]), 1 Japanese database (J-stage), and 7 Korean databases (National Discovery for Science Leaders [NDSL], Korean Medical Database [KMBASE], Oriental Medicine Advanced Searching Integrated System [OASIS] and Korean Traditional Knowledge Portal [KTKP], Korean Studies Information Service System [KISS], Database Periodical Information Academic [DBpia], and the Korean National Assembly Digital Library [KNADL]).

The search terms used were as follows: ((acupuncture, cosmetic OR *miso*-acupuncture OR *miso* acupuncture) AND (prospective OR randomized OR randomized)). The search terms

were slightly modified for acceptable use in each database. Furthermore, the references in searched articles were manually searched to identify further relevant studies.

Study selection

Inclusion criteria

This study included prospective trials that evaluated the effects of acupuncture on cosmetic conditions (e.g., acne vulgaris, facial wrinkles, and anti-aging). We included prospective trials that were either randomized controlled trials (RCTs), non-randomized controlled trials, or prospective, experimental, single-armed studies.

Participants in the included studies underwent acupuncture to improve facial wrinkles, skin conditions, or with a broader goal of anti-aging; obese participants were excluded, regardless of gender, age, or ethnicity.

We included interventions using acupuncture needles inserted into the skin for cosmetic purposes only. Studies that assessed the combined effects of cosmetic acupuncture plus other interventions were also considered when the identical intervention was administered to both the cosmetic acupuncture group and the control group. If a single-armed study used cosmetic acupuncture with other interventions and cosmetic acupuncture was the primary intervention, we included the study after careful review. For control groups, we included any relevant active or sham controls.

We only included relevant outcomes with respect to cosmetic conditions, including total effective rate, wrinkle counts, and any other relevant outcomes measured using devices. Additionally, we included safety reports for the assessment of safety outcomes.

Exclusion criteria

We excluded case reports, non-systematic reviews, surveys, and case series. With respect to cosmetic acupuncture interventions, we excluded auricular acupuncture, embedding acupuncture, or needles attached solely to acupoints. Studies comparing different types of acupuncture were also excluded.

Data extraction

Two independent reviewers (B.C.S. and K.T.L.) screened the titles and abstracts of articles to identify potentially eligible studies and then examined the full texts to assess their final inclusion. Both authors cross-checked eligible studies; any disagreements regarding the extraction of study data were resolved

Table 1. Summary of prospective studies on acupuncture for cosmetic use

Study	Year	Country	Condition	Design	Sample size, female/male	Mean Age (yr), female/male	Treatment Duration (wk)	(A) Acupuncture (B) Control	Outcome measure	Main result	Safety
Huang and Huang [14]	2012	China	Chloasma	RCT	80 (not reported)	33/35	8 wk (20 min/session) 2 times a wk for 4 wk, then one a wk for 4 wk	(A) Cosmetic acupuncture (20 min) + Baiyu power mask (30 min), n=42 -Face Ashi points with ST 7, SI 18, ST2, etc. (B) 0.1% Adapalene gel (daily, for 8 wk), n=38	(1) Total effective rate	(1) 91% (A) vs. 47% (B) (p<0.01)	Not reported
Zhong et al. [8]	2012	China	Acne vulgaris	RCT	70 (67/3)	28.5/28.6	6 wk (20 min/session) 3 times a wk for 6 wk	(A) Cosmetic acupuncture (20 min) + body acupuncture (20 min)+Zhuang herbal medicine mask (30 min), n=35 -Face Ashi points (0.20 x 13 mm needle) -Body acupuncture (LV3, LI4, PC6, SP6, GB41; 0.25 x 30 mm; <i>de-qi</i>) (B) Tanshinone capsule (4 capsules, three times a day, for 6 wk)+ Klein ketone cream (twice a day), n=35	(1) Total effective rate (2) Standard score of Q1 stagnation	(1) 94% (A) vs. 77% (B) (p<0.05) (2) 15.82±5.43 (A) vs. 34.69±8.01 (B) (p<0.01)	(A) 1: Mild itching of the face (B) 2: Dry desquamation, itching
Lee et al. [12]	2010	Korea	Facial wrinkles	Prospective, single-armed, experimental study (before-after study)	6 (6/0)	54.1/-	10 sessions (15 min/session) 3 times a wk	Miso facial acupuncture: Oblique insertion (both) -Sternocleidomastoid m., auricular m., temporalis m., frontal m., masseter m., orbicularis oculi m., risorius m. -0.16 x 30 mm, 0.18 x 40 mm disposable needle	(1) Squares and counts of wrinkles (measured by the facial skin photographing system)	(1) Positive (p<0.05)	Not reported
Kim et al. [10]	2013	Korea	Healthy subjects	Prospective, single-armed, experimental study (before-after study)	10 (10/0)	Not reported	1 session (immediately insertion and removal)	Miso facial acupuncture: 1-3cm depth oblique insertion (both) -Sternocleidomastoid m., auricular m., temporalis m., frontal m., medial pterygoid m., zygomaticus m., masseter m., risorius m., platysma m. -0.16 x 30 mm, 0.18 x 30 mm disposable needle	(1) Facial temperature (2) Blood flow rates	(1) Positive (p<0.05) (2) Positive (p<0.05)	5/10: Non Bruising 2/10: Bleeding 1/10: Bruise+ bleeding Mild adverse event (grade 1)

Table 1. Continued

Study	Year	Country	Condition	Design	Sample size, female/male	Mean Age (yr), female/male	Treatment Duration (wk)	(A) Acupuncture (B) Control	Outcome measure	Main result	Safety
Kwon et al. [11]	2012	Korea	Participants for facial reduction and improve-ment of skin condition	Prospective, single-armed, experimental study (before-after study)	20 (20/0)	Not reported	7 sessions (20 min/ session) 2 times a wk	Miso facial acupuncture Oblique/vertical insertion (both) -Sternocleidomastoid m., auricular m., temporal m., masseter m., digastric m., platysma m., vertical buttress -0.12 x 30 mm, 0.14 x 30 mm, 0.16 x 30 mm disposable needle	(1) Surface distances of face line (measured by 3 dimensional facial scanner) (2) Skin condition (evaluated by water content, trans epidermal water treatment loss (TEWL), erythema index, melanin index)	(1) Positive (p<0.05) (2) Positive (p<0.05)	1 case was dropped out due to pain and swelling
Yun et al. [13]	2013	Korea	Facial wrinkle and aging (Glogau photo aging scale III)	Prospective, open-label, single-armed pilot study (before-after pilot study)	28 (28/0)	50.0/-	5 sessions (10 min/ session) 2 time a wk for the first 2 wk, once a wk for the last wk	Facial cosmetic acupuncture -Temporalis m, epicranial aponeurosis, sternocleidomastoid m., frontalis m., procerus m., corrugators supercilii m., orbicularis oculi m., auricularis m., nasalis m., levator labii superioris m., zygomaticus minor m., zygomaticus major m., orbicularis oris m., risorius m., depressor labii inferioris m., depressor anguli oris m., mentalis m., platysma m. -0.2 x 15 mm, 0.25 x 30 mm disposable needle	(1) Change in the Moire topography (2) A patient-oriented self-assessment of facial elasticity scale	(1) Positive (p<0.05) (2) Positive (p<0.05)	Mild bruising in 14.3% of sessions (20/140 treatment sessions) 1 case dropped out due to pain
Donoyama et al. [9]	2012	Japan	Aging women	Preliminary, prospective, crossover, experimental study	2 (2/0)	50/29	5 sessions (10 min/ session) once a wk	Cosmetic acupuncture: 1-3 mm depth -BL1, GB1, ST1, ST3, ST4, ST7, ST19, CV24, Ex-HN3 and Ex-HN4. -0.10 x 15 mm disposable needle	(1) Water and oil content of facial skin (measured by skin analyzer)	(1) Positive changes compared with no changes in the control condition	Not reported

RCT, randomized controlled clinical trial; m., muscle.

Table 2. Issues related to statistics, ethics, and adherence to reporting guideline on the studies of acupuncture for cosmetic use

Study	Year	Country	Statistics	Ethics	STRICTA reporting guideline item					
					1. Acupuncture rationale	2. Details of needling	3. Treatment regimen	4. Other components of treatment	5. Practitioner background	6. Control or comparator interventions
Huang and Huang [14]	2012	China	Not reported Flaw: no description of statistics	Not reported	Not reported	Partially reported	Reported	Reported	Not reported	Reported
Zhong et al. [8]	2012	China	Chi-square test, t-test Flaw: non	Not reported	Traditional Chinese style (Qi stagnation)	Reported	Reported	Reported	Not reported	Reported
Lee et al. [12]	2010	Korea	Non-parametric test (Wilcoxon signed rank test) Flaw: mean±standard deviation cannot be applied to non-parametric test.	-No IRB approval -Consent (verbal ^{b)} was acquired	Not reported	Partially reported	Reported	Reported	Not reported	NA
Kim et al. [10]	2013	Korea	Parametric test (Student's t-test) Flaw: non-parametric test and median (range) required	-No IRB approval -Written informed consent was acquired	Not reported	Partially reported	Reported	Not reported	Not reported	NA
Kwon et al. [11]	2012	Korea	Parametric (paired t-test) & non-parametric test (Wilcoxon signed rank test) Flaw: all data were summarized as mean±standard deviation, one-tailed test	-No IRB approval -Consent (verbal ^{b)} was acquired	Not reported	Partially reported	Reported	Not reported	Not reported	NA
Yun et al. [13]	2013	Korea	Intension-to-treat analysis Non-parametric test (Wilcoxon signed-rank test) Flaw: mean±standard deviation in non-parametric test	-IRB approval -Written informed consent was acquired	Reported, but insufficient	Reported	Reported	Reported	Not reported	NA
Donoyama et al. [9]	2012	Japan	NA	-Ethic approval was obtained -Written informed consent was acquired	Reported (anatomical base)	Partially reported	Reported	Not reported	Not reported	NA

^{b)}Whether verbal or written informed consent were acquired is unclear.

STRICTA, Standards for Reporting Interventions in Clinical Trials of Acupuncture; IRB, Institutional Review Board; NA, not applicable.

by discussion to reach consensus.

We extracted the following data from the articles selected for inclusion: authors, year of publication, types of cosmetic condition, study design, sample size, interventions, controls, duration of treatment, outcome measures, safety reports, and the main findings (Table 1) [8-14]. We also extracted data regarding statistical procedures, ethics, and adherence to the Standards for Reporting Interventions in Clinical Trials of Acupuncture (STRICTA) (Table 2) [8-15].

Data analyses

All data from outcome measurements were extracted, and we conducted recalculation with respect to effect size (ES) and power (G*power version 3.1.9.2 statistical software) between before and after data by comparing means of matched pairs or goodness-of-fit test by χ^2 test ($\sigma=0.05$, two-tailed). We used mean and standard deviation (or transformed), non-parametric data, or number (%). Initially, we planned to conduct a meta-

analysis. However, high heterogeneity prevented such meta-analysis because different conditions and outcome measures were used.

Results

Study selection

Our search yielded 283 records from English, Chinese, Japanese, and Korean databases, including 52 from the Cochrane library and PubMed, 152 from Korean databases, 61 from the Chinese database, and 18 from the Japanese database. After removal of duplicate studies, 157 records underwent screening of the title and abstract; 145 records were excluded (Fig. 1). The 12 remaining records were analyzed in greater depth to identify relevant studies; 5 were excluded in accordance with our exclusion criteria. Four RCTs were excluded because three trials compared cosmetic acupuncture to different acupuncture styles [16-18], while the other RCT comprised needling of acne




Fig. 1. PRISMA flow diagram of the search and selection process. RCT, randomized controlled trial.

vulgaris [19]; the remaining study constituted a case series, rather than an experimental study [20]. All selection processes are summarized in Fig. 1, in accordance with PRISMA guidelines [7]. Finally, a total of 2 RCTs [8,14] and 5 prospective, experimental studies [9-13] (Chinese, n=2; English, n=2; Korean, n=3) were included in our review. Detailed characteristics of the 7 studies included are summarized in Table 1.

Study characteristics

Two RCTs (n=150) and 5 prospective, single-armed studies (n=66) containing the data of 216 participants were included in the review. The number of participants in the 5 prospective, experimental studies ranged from 2 to 28. All studies were from East Asia: 2 RCTs from China [8,14]; 4 prospective, experimental studies from Korea [10-13]; and 1 from Japan [9]. The number of total sessions was 8.3 ± 5.6 sessions (range, 1-18); the length of each session was 15.8 ± 4.9 minutes (range, 10-20).

The types of conditions were diverse and heterogeneous. The most common conditions treated by cosmetic acupuncture comprised facial care: facial wrinkles (n=2) [12,13], anti-aging (n=1) [9], facial size reduction and skin care (n=1) [11], acne vulgaris (n=1) [8], and chloasma (n=1) [14]; additionally, cosmetic acupuncture was used to perform changes in blood circulation of healthy subjects (n=1) [10] (Table 1).

Outcome measures of the included studies were diverse and heterogeneous (Table 1).

Effects and safety of facial cosmetic acupuncture

All studies were diverse with respect to conditions and outcomes; therefore, we have reported the effects of included studies in a descriptive manner (Table 1).

Effects of facial cosmetic acupuncture on wrinkle care

Two prospective single-armed studies [12,13] assessed the effects of cosmetic acupuncture on squares and counts of wrinkles, or on changes in Moire topography, in facial wrinkle care. Both reported positive effects after 5 or 10 sessions of cosmetic acupuncture treatment (Table 1).

Effects of facial cosmetic acupuncture on skin conditions care

Two prospective experimental studies [9,11] assessed the effects of cosmetic acupuncture on water and oil content, erythema index, and melanin index for the improvement of skin condition. Both reported positive effects (Table 1).

Effects of facial cosmetic acupuncture on other facial conditions

One RCT [14] compared cosmetic acupuncture with 0.1% Adapalene gel for their effects on chloasma by assessing total effective rate; the results of that study were positive ($p < 0.05$). The other RCT [8] tested the effects of cosmetic acupuncture on acne vulgaris, compared to Tanshinone capsule by measuring total effective rate. Its results were also positive ($p < 0.05$).

A prospective study assessed the improvement of blood circulation after cosmetic acupuncture and reported positive results ($p < 0.05$) (Table 1) [10].

Safety reporting

Only 4 of 7 studies [8,10,11,13] reported on the safety of cosmetic acupuncture. Common adverse events were bruising, bleeding, pain, swelling, and itching. All were considered mild adverse events.

Evaluation of statistical methods, ethics, and adherence to acupuncture reporting guidelines

Six studies showed statistical flaws, one did not [8]. The primary shortcoming was the absence of a detailed description of statistical methods; moreover, non-parametric analyses were not used appropriately when the data were not normally distributed or exhibited a small sample size (Table 2).

Two studies reported institutional review board (IRB) approval [9,13], while 5 studies reported obtaining consent from participants; 3 had written informed consent [9,10,13] and 2 simply reported that consent [11,12] had been obtained (Table 2).

Adherence to each item of the STRICTA reporting guidelines was as follows: 3/7 (42.9%) for acupuncture rationale, 7/7 (100%) for details of needling, 7/7 (100%) for treatment regimen, 4/7 (57.1%) for other components of treatment, 0/7 (0%) for practitioner background, and 2/2 (100%) for control or comparator interventions (Table 2).

Recalculation of effect size and power

Two RCTs showed $ES=1.43$, $power=1.0$ [14] and $ES=0.72$, $power=0.99$ [8], when recalculated by χ^2 test.

Prospective experimental studies showed $ES=0.53$, $power=0.19$ with respect to counts of skin furrows [12]; $ES=1$, $power=0.80$ with respect to changes in facial temperature [10]; $ES=1.87$, $power=1.0$ with respect to water count [11]; and $ES=0.60$, $power=0.86$ with respect to mean change in Moire topography [13]. One study could not be recalculated because

minimal data were reported [9].

Risk of bias of RCTs

One RCT reported use of a random number table [8] but the other RCT did not report any use of randomization [14]. Blinding of practitioner, patients, and assessor was not reported in either RCT [8,14].

Discussion

The purpose of our systematic review was to explore the current status of facial cosmetic acupuncture and to evaluate the effects and safety of facial acupuncture for cosmetic use. Our primary finding was that many studies used defective research methodology and exhibited poor adherence to the STRICTA reporting guidelines for clinical acupuncture trials.

Of the 2 RCTs from China, only 1 reported an adequate randomization method (random number table) [8]. Five prospective studies also exhibited a high risk of bias because they were not controlled studies [21]. No trial adopted a meaningful blinding method for practitioner, patients, and assessor. To explore the impact of cosmetic acupuncture with respect to any facial conditions, we conducted recalculation of ES and power of the included studies. However, 1 prospective study [12] did not exhibit full power (0.19); this indicated an underpowered study with potentially unreliable results. Future trials should include sufficient sample sizes; pre-enrollment sample size calculation is highly recommended. Another concern involves the research design of the included studies. Most included studies adopted heterogeneous definitions of wrinkles, skin care, anti-aging, acne vulgaris, and chloasma; thus, our review could not draw any meaningful conclusions or generate a combined conclusion from the synthesis of included studies.

An important finding is that treatment trends differ widely between China, Korea, and Japan. Generally, Korean acupuncture used *Miso* facial acupuncture for the cosmetic treatment of face and neck muscle, without using meridian or acupuncture theory [10-13]; Chinese and Japanese acupuncture comprised face *Ashi* points with meridian theory. Additionally, Chinese acupuncture used moderate needle length [8,14], whereas the Japanese style used shallow needling to the face (1-3 mm depth) [9].

Our analysis had several limitations. Although we found 5 Chinese RCTs, we excluded 3 of those [16-18] because they compared cosmetic acupuncture with different types of body acupuncture; thus, it was not possible to clearly elucidate acu-

puncture effects. Another concern is the small sample size of the included studies. Our recalculations showed moderate ES and power; however, this may be a result of a biased, exaggerated effect [22]. All RCTs and prospective studies included in this study presented methodological weaknesses; this prevented us from making clear conclusions regarding the efficacy of cosmetic acupuncture in facial care.

One article reported extensive facial sclerosing lipogranulomatosis as a complication of cosmetic acupuncture [23]. However, overall adverse events of cosmetic acupuncture reported in our review appear tolerable.

Future trials should consider standardized cosmetic acupuncture intervention and comparison with acceptable controls, such as sham or active standard treatments with a standardized regimen. Additionally, researchers should adhere to STRICTA reporting guidelines with respect to acupuncture rationale (42.9%), other components of treatment (57.1%), and practitioner background (0%) [15].

Conclusion

Our systematic review of 2 RCTs and 5 prospective, single-armed studies demonstrated that there are a few trials with high-quality evidence regarding cosmetic acupuncture for facial cosmetic conditions. The safety of cosmetic acupuncture appears tolerable; however, well-designed high-quality studies, such as sham-controlled studies with standardized cosmetic acupuncture interventions, are needed to determine its effects on facial cosmetic conditions.

Acknowledgments

This study was supported by clinical research grant from Pusan National University Hospital in 2018.

Conflicts of interest

The authors have nothing to disclose.

References

1. International Society of Aesthetic Plastic Surgery (ISAPS). ISAPS international survey on aesthetic/cosmetic procedures performed in 2016 [Internet]. Hanover (NH): ISAPS, 2014 [cited 2018 Aug 24]. Available from: <https://www.isaps.org/blog/isaps-statistics/>.

2. Maisel A, Waldman A, Furlan K, Weil A, Sacotte K, Lazaroff JM, et al. Self-reported patient motivations for seeking cosmetic procedures. *JAMA Dermatol* 2018;154:1167-74.
3. Barrett JB. Acupuncture and facial rejuvenation. *Aesthet Surg J* 2005;25:419-24.
4. Doran V. An introduction to facial revitalisation acupuncture. *Eur J Orient Med* 2007;5:4-9.
5. Kim TY. Trend analysis of facial cosmetic acupuncture study based on the Korean Traditional Medicine. *J Acupunct Res* 2013;30:125-37.
6. Fang Y, Fumihiko S, Mamoru U, Masahiro W, Akio T. Cosmetic acupuncture for Oriental Medicine treatment. *J Osaka Dent Univ* 2012;46:5-9.
7. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
8. Zhong J, Fu LL, Fang G, Zhang XP, Lin C. Clinical study on the treatment of Qi stagnation type acne vulgaris with acupuncture combined with Zhuang herbal medicine mask. *Nat Med Front China* 2012;7:55-6.
9. Donoyama N, Kojima A, Suoh S, Ohkoshi N. Cosmetic acupuncture to enhance facial skin appearance: a preliminary study. *Acupunct Med* 2012;30:152-3.
10. Kim TY, Bak JP, Kim YM. Changes of facial temperature and blood flow rates by treatment of Miso facial rejuvenation acupuncture. *Korean J Orient Physiol Pathol* 2013;27:481-6.
11. Kwon GS, Kim JH, Lee KA, Lee SJ, Song JH, Song CH, et al. The effect of Miso facial acupuncture on facial reduction and improvement of skin condition. *J Korean Acupunct Moxibustion Med Soc* 2012;29:7-18.
12. Lee KM, Lim SC, Kim JS, Lee BH. A clinical study on facial wrinkles treated with Miso facial acupuncture-measured by the facial skin photographing system-. *J Korean Acupunct Moxibustion* 2010;27:101-7.
13. Yun Y, Kim S, Kim M, Kim K, Park JS, Choi I. Effect of facial cosmetic acupuncture on facial elasticity: an open-label, single-arm pilot study. *Evid Based Complement Alternat Med* 2013;2013:424313.
14. Huang YS, Huang W. Nursing experience of treatment of chloasma with cosmetic acupuncture combined with Biayu powder mask. *Mod J Integr Tradit Chin West Med* 2012;21:2945-6.
15. MacPherson H, Altman DG, Hammerschlag R, Youping L, Taixiang W, White A, et al; STRICTA Revision Group. Revised STandards for Reporting Interventions in Clinical Trials of Acupuncture (STRICTA): extending the CONSORT statement. *PLoS Med* 2010;7:e1000261.
16. Chang C, Lyu Y, Cao Q, Zheng B, Wan Y, Xu J. Comparative study on repairing effect of lance needle and cosmetic needle for pustular acne. *Acta Univ Tradit Med Sin Pharmacol Shanghai* 2013;27:51-53, 57.
17. Chen J. [Clinical observation of acupuncture treatment for 68 Chloasma patients]. *Shenzhen J Integr Tradit Chin West Med* 2015;(14):61-2. Chinese.
18. Zhang J, Lu Y, Liu Q, Cao Q, Jiang Y, Ding J, et al. Therapeutic observation on lance needle for acne vulgaris. *Shanghai J Acupunct Moxibustion* 2013;(11):929-30.
19. Zeng X, Yu G. [75 Cases of acne vulgaris treated by staged nursing care]. *Chin J Ethnomed Ethnopharm* 2012;(16):106-7. Chinese.
20. Kwon NH, Kim CY, Shin YJ, Seo S, Song JH, Baek YH, et al. Clinical study on facial skin furrow measurement changes after Miso facial rejuvenation acupuncture. *J Korean Acupunct Moxibustion Soc* 2009;26:133-40.
21. Gluud LL. Bias in clinical intervention research. *Am J Epidemiol* 2006;163:493-501.
22. Button KS, Ioannidis JP, Mokrysz C, Nosek BA, Flint J, Robinson ES, et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013;14:365-76.
23. Bashey S, Lee DS, Kim G. Extensive facial sclerosing lipogranulomatosis as a complication of cosmetic acupuncture. *Dermatol Surg* 2015;41:513-6.